DISCUSSION PAPERS IN ECONOMICS

Working Paper No. 16-07

Oracle Efficiency in Additive Partially Linear Triangular Systems

Eric Penner
University of Colorado Boulder

October 2016

Department of Economics

University of Colorado Boulder
Boulder, Colorado 80309

© October 2016 Eric Penner

1 Introduction

The treatment of endogeneity in semiparametric models has been the subject of intense interest in recent years. In particular, since Newey et al. (1999) in which the control function approach to endogeneity in nonparametric models is established, triangular systems of equations have garnered a great deal of attention see, inter alia, Geng et al. (2016), Ozabaci et al. (2014), Su and Ullah (2008), and Martins-Filho and Yao (2012). The value of the control function approach is clear, providing a method of dealing with endogeneity that minimizes the risk of model misspeci cation and is easy to implement. I use a control function to handle endogeneity in a partially linear model. Accordingly, this model consists of D+1 equations; a single partially linear primary equation (1) with a mixture of endogenous and exogenous regressors, as well as D fully nonparametric secondary equations (2) having only exogenous regressors. Thus, consider the following triangular system of equations where a set of exogenous regressors enters the primary equation (1) parametrically while allowing endogenous regressors to enter nonparametrically.

$$Y = {}_0 + Z^0 {}_1 + h(X) + "; \tag{1}$$

$$X = m(W) + V; \tag{2}$$

$$E(VjW) = 0; \quad E("jW; X) = E("jV): \tag{3}$$

Y is a scalar random variable, X is a D dimensional vector of endogenous random variables in that $E("jZ; X) = E("jX) \neq 0$. W is a q dimensional random vector, and Z is a $p < q$ dimensional subvector of W. " and V are random disturbances. $[{}_0 \quad {}^0_1]^0 \in R^{p+1}$ and $h(\,)$ are unknown parameters of interest. $m(W) : R^q \to R^D$ is a vector of D real valued functions $m_d(W)$ where $d \in \{1; 2; : : : ; D\}$.

Specifying a partially linear form for the primary equation has the distinct advantage of allowing one to impose a parametric form for some regressors, when justi ed, while allowing others to be the arguments of a much broader class of functions. Additionally, Robinson (1988) showed that in models consisting solely of equation (1) and where $E("jZ; X) = 0$, his estimator for $[{}_0 \quad {}^0_1]^0$, the nite dimensional parameter of central interest, is consistent and $\sqrt{n}$ asymptotically normal. Furthermore the asymptotic covariance matrix of the estimator for $[{}_0 \quad {}^0_1]^0$

2 Moment Conditions, Identification, and Estimation

2.1 Moment Conditions

$E("jX) \neq 0$ implies that $"$ is not orthogonal to the space of square integrable functions of X. The approach taken to deal with this lack of orthogonality is a variation on the control function approach of Newey et al. (1999). In particular, defining $u = " \quad f(V)$, I assume,

$$E["jZ; X; V] =$$

where,

(Z_c), and $_Y$ are estimated with $\hat{}_{Z_c} = n^{-1}\sum_{i=1}^{n} \hat{}_i Z_{ci}$ and $\hat{}_Y = n^{-1}\sum_{i=1}^{n} \hat{}_i Y_i$ so that,

$$\hat{H}(Z_{ci}) = \sum_{d=1}^{p} [\hat{H}_1^d(Z_{ci}) + \hat{H}_2^d(Z_{ci})]; \qquad \hat{H}(Y_i) = \sum_{d=1}^{p} [\hat{H}_1^d(Y_i) + \hat{H}_2^d(Y_i)]; \tag{25}$$

$$\hat{H}(Z_{ci}) = \hat{H}(Z_{ci}) \quad (2D \quad 1)\hat{}_{Z_c}; \quad \hat{H}(Y_i) = \hat{H}(Y_i) \quad (2D \quad 1)\hat{}_Y : \tag{26}$$

Let $\hat{H}_n(Y) = [\ \hat{H}(Y_1) \quad \hat{H}(Y_2) \quad \hat{H}(Y_n)\ ]^0$, $\hat{H}_n(Z_c) = [\ \hat{H}(Z_{c1}) \quad \hat{H}(Z_{c2}) \quad \hat{H}(Z_{cn})\ ]^0$ and $\hat{H}_n(Z) = [\ \hat{H}_n(Z_1) \quad \hat{H}_n(Z_2) \quad \hat{H}_n(Z_p)\ ]$ where $Y_n = [\ Y_1 \quad Y_2 \quad Y_n\ ]^0$, and $Z_n = [\ Z_1 \quad Z_2 \quad Z_n\ ]^0$ and let,

$$\hat{}_{ci} = Z_{ci} \quad \hat{H}(Z_{ci}); \qquad \hat{}_i = Z_i \quad \hat{H}(Z_i); \qquad \hat{}_n = Z_n \quad \hat{H}_n(Z): \tag{27}$$

Step Four: Estimation of $(\ _0 \quad {}_1^0)^0 \in R^{p+1}$.
I regress $\hat{}_n^{1=2}(Y_n \quad \hat{H}_n(Y))$ on $\hat{}_n^{1=2}\hat{}_n = \hat{}_n^{1=2}(Z_n \quad \hat{H}_n(Z))$ and estimate $_1$ as,

$$\hat{}_1 = [\hat{}_n^0 \hat{}_n \hat{}_n]^{-1}\hat{}_n^0 \hat{}_n(Y_n \quad \hat{H}_n(Y)): \tag{28}$$

where $\hat{}_n \quad \hat{}_n(X; \hat{V}) = \mathrm{diag}(f\hat{}(X_i; \hat{V}_i)g_{i=1}^n)$, Lastly $_0$ is estimated by,

$$\hat{}_0 = \hat{}_Y \quad \hat{}_Z^0 \hat{}_1: \tag{29}$$

3 Asymptotic Results

In this section, I study the asymptotic properties of $[\hat{}_0 \ \hat{}_1^0]^0$. To do so, I rst de ne vector and matrix norms that I will use throughout this paper. For $m; n \in \mathbb{N}$, let A be a matrix of order $(m \quad n)$, r be a random vector of order $(m \quad 1)$ whose elements have nite second moments, and be a non stochastic vector of order $(m \quad 1)$. De ne $\|A\| = \sqrt{\mathrm{trace}(AA')}$, the Frobenius matrix norm, de ne the matrix spectral norm $\|A\|_{sp} = \max_{1 \ i \ m} {}_i(AA')^{1=2}$ where $f\ _i(AA')g_{i=1}^m$ the collection of all eigenvalues of AA'. De ne the inner product $\|r\|_2^2 = E(r'r)$, and lastly de ne $\|c\|_E = (c'c)^{1=2}$ the typical Euclidean norm in nite dimensional space R^m.

3.1 Assumptions

Assumption A1: The B-spline basis functions used in step one satisfy the following: (i) Let $f l_n g_{n=1}^1$ be a nondecreasing sequence of natural numbers such that $l_n^2=n = o(1)$. Let $G_{W_a} \quad R$ be the compact support of W_a, and without loss of generality. let $G_{W_a} = [0; 1]$. For some $k \quad 3$ let $f t_j g_{j=1}^{l_n+2\ k}$ be a knot

set ^{2} for G_{W_a}. Now de ne the following functions

$$b_{j;1}(W_a) = \begin{cases} 1 & \text{if } W_a \ 2 \ [t_j ; t_{j+1}) \\ 0 & \text{if } W_a \ 2 \not\in [t_j ; t_{j+1}); \end{cases} \qquad \text{and} \qquad !_{j;k}(W_a) = \frac{W_a \quad t_j}{t_{j+k \ 1} \quad t_j}:$$

Now, de ne the normalized B-spline basis of order $k \ 2$ N functions recursively,

$$b_{j;k}(W_a) = !_{j;k}(W_a) b_{j;k \ 1}(W_a) + (1 \quad !_{j+1;k}(W_a)) b_{j+1;k \ 1}(W_a) \qquad \text{and} \qquad B_{j;k}(W_a) = \frac{b_{j;k}(W_a)}{jj b_{j;k}(W_a) jj_2}$$

where $\{ B_{j;k}(W_a) \}_{j=1}^{I_n + 2 \ k}$ is a sequence of uniformly bounded, positive, twice integrable, compactly supported B-spline basis functions with which I construct,

$$B_n(W_{ai})' = \quad B_{1;k}(W_{ai}) \qquad B_{I_n + 2 \ k;k}(W_{ai})$$

(ii) De ne $Q_{BB} = E[B_n(W_i) B_n(W_i)']$, $Q_{nBB} = n^{\ 1} \sum_{i=1}^{n} B_n(W_i) B_n(W_i)'$, $Q^d_{BBV} = E\left[B_n(W_i) B_n(W_i)' V_{di}^2 \right]$, and $Q^d_{nBBV} = n^{\ 1} \sum_{i=1}^{n} B_n(W_i) B_n(W_i)' V_{di}^2$. Assume that for n su ciently large, there exist constants $c_{Bl} ; c_{Bh} ; c_{BV} \ 2 \ R^+$ such that

$$0 < c_{Bl} \quad \min(Q_{BB}) \quad \max(Q_{BB}) \quad c_{Bh} < 1 \qquad \text{and} \qquad 0 < \quad \max(Q^d_{BBV}) \quad c_{BV} < 1$$

Remarks $^{n+2 \ k}_{z}$ $^{d}_{R}$

arbitrarily small " > 0

$$_0 = \inf\ f x\ 2\ N : x >\ 1=4 f\ g;$$

$$_1 = \inf\ f x\ 2\ N : x >\ 1=4 e g;$$

$$_2 = \inf\ f x\ 2\ N : x >\ 1=4 b g;$$

$$_3 = \inf\ \ x\ 2\ N : x >\ \ 4\ \ \min\ \ \frac{1}{4(D+1)}; \frac{2k\ \ 1}{4k(D+1)}; \frac{3k\ \ 3}{16k}\ \ \ \ " \ \ \ \ _1) \ :$$

Remarks : Assumptions (i) - (vi), are su cient conditions for the rates of convergence derived in Lemma 3. Note that the conditions on h_0

and M_{2n} dominate L_n the rate of convergence derived in Theorem 1 is the same as if $m(W)$ is a known function. Furthermore the same holds true for the uniform rate of convergence of density ratios $\hat{\ }_i$, $\hat{\ }^d_{1i}$, and $\hat{\ }^d_{2i}$.

Theorem 1. Under Assumptions A1 - A5,

$$\sup_{X_{di} \in G_{X_d}} |\hat{p}(X_{di}) - p(X_{di})| = O_p(M_{1n}); \qquad \sup_{V_{di} \in G_{V_d}} |\hat{p}(\hat{V}_{di}) - p(V_{di})| = O_p(L_n + M_{1n});$$

$$\sup_{X_i, V_i \in G_{X,V}} |\hat{p}(X_i, \hat{V}_i) - p(X_i, V_i)| = O_p(L_n + M_{2n}); \qquad \sup_{X_i, V_i \in G_{X,V}} |\hat{\ }(X_i, \hat{V}_i) - (X_i, V_i)| = O_p(L_{0n});$$

$$\sup_{X_i, V_i \in G_{X,V}} |\hat{\ }(X_{di}, \hat{V}_i) - (X_{di}, V_i)| = O_p(L_{0n}); \qquad \sup_{X_i, V_i \in G_{X,V}} |\hat{\ }(X_i, \hat{V}_{di}) - (X_i, V_{di})| = O_p(L_{0n}).$$

Theorem 2. Under assumptions A1 - A6, $\hat{\ }_Y - {}_Y = O_p(L_{0n})$, $\hat{\ }(Z_c) - (Z_c) = O_p(L_{0n})$ and

$$\sup_{X_{di} \in G_{X_d}} |\hat{H}^d_1(Y_i) - H^d_1(Y_i)| = O_p(L_{1n}); \qquad \sup_{V_{di} \in G_{V_d}} |\hat{H}^d_2(Y_i) - H^d_2(Y_i)| = O_p(L_{2n});$$

$$\sup_{X_{di} \in G_{X_d}} |\hat{H}^d_1(Z_{ci}) - H^d_1(Z_{ci})| = O_p(L_{1n}); \qquad \sup_{V_{di} \in G_{V_d}} |\hat{H}^d_2(Z_{ci}) - H^d_2(Z_{ci})| = O_p(L_{2n});$$

$$\sup_{X_i, V_i \in G_{X,V}} |\hat{H}(Y_i) - H(Y_i)| = O_p(L_n); \qquad \sup_{X_i, V_i \in G_{X,V}} |\hat{H}(Z_{ci}) - H(Z_{ci})| = O_p(L_n).$$

Remarks : The uniform rate of convergence of these estimators are the sum of the uniform rates of convergence of $\hat{m}_d(W_{ai})$, $\hat{\ }(X_{di}, \hat{V}_i)$ or $\hat{\ }(X_i, \hat{V}_{di})$, and a typical NW estimator, in each case except for those terms where the conditional expectation is taken with respect to (w.r.t) V. In this case there is an additional term b_2^{-1}

exercise are summarized in Table 1, which lists the bias (B), the standard deviation (S), the root mean squared error (R), and the median squared error (D), for each estimator. Also listed are the mean squared error (M1), and (M2) for the estimation of $m_1(W)$ and $m_2(W)$ respectively. Note, that the Table and Figures summarize only those estimated values within the 2% - 98% sample quantile range. Also note that M1 and M2 values for $\hat{\ }_{1;M}$ are left blank as in this case $m_1(W)$ and $m_2(W)$ are taken to be known functions. In addition to the results summarized in Table 1, Rosenblatt kernel density estimates for the absolute error of all three estimators are shown in Figure 1 for a sample size of $n = 200$, and in Figure 2 for a sample size of $n = 400$.

The additive model presented in the paper is a restriction on the model developed in Geng et al. (2016) who assume no additivity. As a result one should expect that the estimator in this paper will have better nite sample performance for two reasons. Firstly, since both primary equations in this study are additive functions, the estimator in this paper should perform better than Geng et al. (2016) regardless of the degree of endogeneity or sample size as it is has a sharper speci cation. Secondly, as discussed in the introduction, using an additive B spline estimator for $m(W)$ in this setting avoids the \curse of dimensionality" relative to estimators that do not exploit additivity. Consequently, the estimators for $m_1(W)$ and $m_2(W)$ in this paper should outperform Geng et al. (2016) who use Nadaraya Watson estimators. Table 1 demonstrates that this is true, as in all cases the mean squared error for $\hat{m}(W)$ is smaller than $\hat{m}_G(W)$.

The structure of this Monte Carlo exercise is designed to demonstrate changes in the nite sample properties of each estimator due to variations in data generating process, sample size, and degree of endogeneity. As for variations in sample size, in Table 1 it is clear that increasing the sample size has the expected e ect since, in all cases, the bias, standard deviation, and mean squared error of each estimator decline, supporting the consistency results derived for each. As the level of endogeneity increases one should expect that the two disadvantages inherent in Geng et al. (2016) discussed above should be compounded since comparatively poorer estimates $\hat{m}_G(W)$, due to a more general speci cation for $m(W)$, are used in a second stage estimator

^

Table 1: Finite Sample Performance

	$_e$ = 0:3						$_e$ = 0:6						$_e$ = 0:9					
	B	S	R	D	M1	M2	B	S	R	D	M1	M2	B	S	R	D	M1	M2
DGP=1	n=200																	
$\hat{}_1$	-0.5982	0.1075	0.3694	0.3618	0.2938	0.3066	-0.6400	0.1016	0.4199	0.4104	0.2998	0.3102	-0.6704	0.0967	0.4587	0.4516	0.2980	0.3089
$\hat{}_{1;G}$	-0.6641	0.1487	0.4631	0.4322	0.7041	0.6926	-0.8065	0.1362	0.6690	0.6465	0.7046	0.6905	-0.8787	0.1112	0.7844	0.7747	0.6997	0.6964
$\hat{}_{1;M}$	1																	



Figure 1: Estimated Densities for Absolute Error of estimators of $_1$, n = 200



Figure 2: Estimated Densities for Absolute Error of estimators of $_1$, n = 400

Recall that the sole di erence between estimators $\hat{\theta}_1$ and $\hat{\theta}_{1;M}$ is whether $m(W)$ is a known or estimated. In Theorem 3, I have shown that the asymptotic variance of $\hat{\theta}_1$ is identical to $\hat{\theta}_{1;M}$, a property which I call Oracle e ciency in the somewhat atypical sense that I describe in the introduction. The purpose of comparing $\hat{\theta}_1$ and $\hat{\theta}_{1;M}$ here is to determine to what extent this Oracle property is evident in nite samples. The ndings presented in Table 1, and both Figures reinforce the Oracle e ciency

$$= h_d(X_d) \int \frac{g(X;V)}{p(X;V)} \frac{p(X;V)}{p(X_d)} d(X_{-d};V) + \int \left(\sum_{j \neq d}^{D} h_j(X_j) + \sum_{j=1}^{D} f_j(V_j) \right) \frac{g(X;V)}{p(X;V)} \frac{p(X;V)}{p(X_d)} d(X_{-d};V)$$

$$= h_d(X_d) \int g(X_{-d};V) d(X_{-d};V) + \sum_{j \neq d}^{D} \int h_j(X_j) g(X_{-d};V) d(X_{-d};V) + \sum_{j=1}^{D} \int f_j(V_j) g(X_{-d};V) d(X_{-d};V)$$

$$= h_d(X_d) + \sum_{j \neq d}^{D} E[h_j(X_j)] + \sum_{j=1}^{D} E[f_j(V_j)] = h_d(X_d):$$

In a similar manner, $E\left[\sum_{j=1}^{D} h_j(X_j) + \sum_{j=1}^{D} f_j(V_j) \mid V_d \right] = f_d(V_d)$. Additionally note that,

$$E[(Y - Z\beta_1)] = \int (Y - Z\beta_1)$$

$$= \beta_0 E[1] + \sum_{d=1}^{D} E[h_d(X_j)] + \sum_{d=1}^{D} E[f_d(V_j)] + E[\varepsilon]$$

$$= \beta_0 \int \frac{g(X;V)}{p(X;V)} p(X;V) dX dV + \sum_{d=1}^{D} \int h_d(X_d) \frac{g(X;V)}{p(X;V)} p(X;V) dX dV$$

$$+ \sum_{d=1}^{D} \int f_d(V_d) \frac{g(X;V)}{p(X;V)} p(X;V) dX dV + E[E(\varepsilon | Z;X;V)]$$

$$= \beta_0 + \sum_{d=1}^{D} E[h_d(X_d)] + \sum_{d=1}^{D} E[f_d(V_d)]$$

$$= \beta_0:$$

As a result, provided β_Y and β_Z exist, $\beta_1 = 0$ Td [(j)]TJ/F11 9.9626 Tf 2.767 0 Td [(Z)-16(;)-167(X)-232.767 0 Td [(h)]TJ/F10 6.9l(ti)1[(d46.496)]TJwher

$$= {}_0 + (2D \quad 1)\ {}_Y \quad (2D \quad 1)\ \frac{{}_0}{Z}\ {}_1 + H(Z)^0\ {}_1 + h(X) + f(V) \quad (2D \quad 1)\ {}_Y$$
$$= {}_0 + H\ (Z)^0\ {}_1 + h(X) + f(V):$$

$$= \sum_{j=1}^{p} \int \frac{g(X_{-d}, V)}{p(X, V)} E[A \mid V_j] p(X, V)\, dX\, dV + (2D - 1) E[A] E[\mid X_d]$$

$$= H_1^d(A) - H_1^d(A) \sum_{j \neq d} \int E[A \mid X_j] p(X_j)\, dX_j \sum_{j=1}^{p} \int E[A \mid V_j] p(V_j)\, dV_j + (2D - 1) E[A]$$

$$= (2D - 1) E[A] \sum_{j \neq d} E[A] \sum_{j=1}^{p} E[A] = 0.$$

Similarly, $E[(A - H(A)) \mid V_d] = 0$. □

Lemma 3. Under Assumption A2

i) $l_n^{\,k} = o(n^{-1/2})$,

ii) $l_n = [\sqrt{n} b_2] = o(n^{-1/4})$,

iii) $l_n = [n(h_1^D h^{D=}$

By L'Hôpital's Rule.

Part xv) : By xii) and xiii), $M_{1n} = \left[\frac{\log(n)}{nh_0}\right]^{1=2} + h_0{}^0 = o(n^{1=4}) + o(n^{1=4})$.

Part xvi) : By A6: $_3 > 1=4c$ let $_3 = 1=4c + "$,

$$h_1{}^3 n^{1=4} + h_2{}^3 n^{1=4} \quad n^{c\ 3} n^{1=4} = n^{c(1=4c+")+1=4} = n^{c"} = o(1):$$

Part xvii) : By A6: $c < 1=4(D+1)$ let $c = 1=4(D+1) \quad "$,

$$\log(n) n^{1}h_1{}^{D} h_2{}^{(D+2)} n^{1=2} \quad \log(n) n^{1=2} n^{2(D+1)c} = \log(n) n^{1=2+2(D+1)(1=4(D+1)\quad ")}$$
$$= n^{1=2+1=2\quad 2(D+1)"} = n^{2(D+1)"} = o(1):$$

Part xviii) : By (xiv) and (xv), $M_{2n} = \left[\frac{\log(n)}{nh_1^D h_2^D}\right]^{1=2} + h_1{}^3 + h_2{}^3 = o(n^{1=4}) + o(n^{1=4})$.

Part xix) : By A6: $a > 1=2k$, and $c < (2k \quad 1)=[4k(D+1)]$, let $a = 1=2k + "$ and $c = (2k \quad 1)=[4k(D+1)] \quad "$,

$$l_n n^{1}h_1{}^{D} h_2{}^{(D+2)} \quad n^a n^{1} n^{2c(D+1)}$$
$$= n^{1=2k+"\quad 1+2(D+1)(2k\quad 1)=[4k(D+1)]\quad 2(D+1)"}$$
$$= n^{1=2k\quad 1+1\quad 1=2k\quad (2D+1)"}$$
$$= n^{(2D+1)"} = o(1):$$

Parts xx) - xxv) : Proofs follow, mutatis mutandis, from parts xix) - xxiv) are mutatis mutandis virtually identical to the proofs of parts xii) - xiv) please consult them.

Parts xxvi) & xxvii) By the above results,

$M_n = M_{1n} + M_{2n} = o(n^{1=4})$; $L_{0n} = L_n + M_n = o(n^{1=4})$; $L_{1n} = L_n + M_n + N_{1n} = o(n^{1=4})$;

$L_{1n} = L_n + M_n + N_{2n} + b_2{}^1 \left[\frac{l_n}{n}\right]^k + l_n{}^k = o(n^{1=4})$; $L_n = L_{1n} + L_{2n} = o(n^{1=4})$:

□

if $\lambda_{min}(A) \geq 0$ then $\sigma_{min}(A) = \lambda_{max}(A) = \|A\|_2 = \sigma_{max}(A)$:

$$= \min(Q_{BB}) - o_p(1):$$

Combining the previous two statement gives (i),

$$\min(Q_{BB}) + o_p(1) \geq \min(Q_{nBB}) \geq \min(Q_{BB}) - o_p(1):$$

Part (iii.) Let h = 1 so that $_h(\cdot) = \max(\cdot)$.

$$\begin{aligned}\max(Q_{nBB}) &= \max(Q_{BB} + Q_{nBB} - Q_{BB}) \\ &\leq \max(Q_{BB}) + \max(Q_{nBB} - Q_{BB}) \\ &\leq \max(Q_{BB}) + \|Q_{nBB} - Q_{BB}\| \\ &= \max(Q_{BB}) + o_p(1):\end{aligned}$$

Furthermore,

$$\begin{aligned}\max(Q_{nBB}) &= \max(Q_{BB} + Q_{nBB} - Q_{BB}) \\ &\geq \max(Q_{BB}) + \max(Q_{nBB} - Q_{BB}) \\ &\geq \max(Q_{BB}) + \|Q_{nBB} - Q_{BB}\| \\ &= \max(Q_{BB}) + o_p(1):\end{aligned}$$

Combining the previous two statements gives (ii)

$$\max(Q_{BB}) + o_p(1) \geq \max(Q_{nBB}) \geq \max(Q_{BB}) - o_p(1):$$

Part (iv)

$$= (h_1^D h_2^D)^{-1} \int D_d K_3[H^{-1} f(X_j^0, V_j^0)' - (X_i^0, V_i^0)']g] p(X_j, V_j, W_j) p(W_j)^{-1} dX_j\, dV_j$$

$$= (h_1^D h_2^D)^{-1} \int D_d K_3[(\epsilon_1, \epsilon_2)]^2 p(X_i + \epsilon_1 h_1, V_i + h_2 \epsilon_2, W_j) p(W_j)^{-1} h_1^D d\epsilon_1 h_2^D d\epsilon_2$$

$$\sup_{X,V,W \in G_{XVW}} p(X,V,W) \inf_{W \in G_W} p(W)^{-1} \int D_d K_3[(\epsilon_1^0, \epsilon_2^0)']^2 d\epsilon_1 d\epsilon_2 = O(1).$$

Consequently, uniformly in $G_{X,V}$,

$$E\left[(h_1^D h_2^D)^{-1} D_d K_{3ji}^2(W_j, S_i)\right] = O(1).$$

By Assumption A2, $\lim_{\epsilon_1, \epsilon_2 \to 1} K_3[(\epsilon_1^0, \epsilon_2^0)] - K[(\epsilon_1, \epsilon_2)] = 0$ then by A2 and A4

$$E\left[(h_1^D h_2^{D+1})^{-1} D_d K_{3ji}(W_j, S_i)\right] = E\left[(h_1^D h_2^{D+1})^{-1} D_d K_3[H^{-1} f(X_j^0, V_j^0)' - (X_i^0, V_i^0)']g](W_j, S_i)\right]$$

$$= (h_1^D h_2^{D+1})^{-1} \int D_d K_3[H^{-1} f(X_j^0, V_j^0)' - (X_i^0, V_i^0)']g] p(X_j, V_j, W_j) p(W_j)^{-1} dX_j\, dV_j$$

$$(h_1^D h_2^{D+1})^{-1} \left[K_3[H^{-1} f(X_j^0, V_j^0)' - (X_i^0, V_i^0)']g] p(X_j, V_j, W_j) p(W_j)^{-1}\right]_1^1$$

$$+ (h_1^D h_2^{D+1})^{-1} \int h_2 K_3[H^{-1} f(X_j^0, V_j^0)' - (X_i^0, V_i^0)']g] D_d p(X_j, V_j, W_j) p(W_j)^{-1} dX_j\, dV_j$$

$$\sup_{X,V,W \in G_{XVW}} p(X,V,W) \inf_{W \in G_W} p(W)^{-1} (h_1^D h_2^{D+1})^{-1} K_3[(1,1)] - K_3[(\epsilon_1, 1)]$$

$$+ (h_1^D h_2^D)^{-1} \int K_3[(\epsilon_1^0, \epsilon_2^0)'] D_d p(X_i + h_1 \epsilon_1, V_i + h_2 \epsilon_2, W_j) p(W_j)^{-1} h_1^D d\epsilon_1 h_2^D d\epsilon_2$$

$$\sup_{X,V,W \in G_{XVW}} D_d p(X,V,W) \inf_{W \in G_W} p(W)^{-1} \int |K_3[(\epsilon_1^0, \epsilon_2^0)']| d\epsilon_1 d\epsilon_2 = O(1)$$

Consequently, uniformly in $G_{X,V}$,

$$E\left[(h_1^D h_2^{D+1})^{-1} D_d K_{3ji}(W_j, S_i)\right] = O(1).$$

Note that,

$$E(B_n(W_i)' B_n(W_i)) = \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} E\left[B_{J;k}(W_{ai})^2\right] = \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} E\left[\frac{b_{J;k}(W_a)^2}{\|b_{J;k}(W_a)\|_2^2}\right] = \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} 1 = q(l_n + 2k).$$

Let $|B_n(W)|$ be an element wise absolute value. For $i \neq j$ note that,

$$E\left[|B_n(W_i)'| |B_n(W_j)|\right] = \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} E\left(|B_{J;k}(W_{ai})|\right) E\left(|B_{J;k}(W_{aj})|\right) = \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} E\left(|B_{J;k}(W_{ai})|\right)^2$$

$$= \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} (\|b_{J;k}(W_{ai})\|_2^2)^{-1} \left[\int_{t_J}^{t_{J+k}} b_{J;k}(W_{ai}) p(W_{ai}) dW_{ai}\right]^2$$

$$= \max_{1 \le a \le k} \max_{1 \le J \le l_n + 2k} (\|b_{J;k}(W_{ai})\|_2)^{-2} \sup_{W_a \in G_{W_a}} |b_{J;k}(W_a)|^2 |p(W_a)|^2 \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} \left[\int_{t_J}^{t_{J+k}} dW_{ai}\right]^2$$

$$= O(l_n) \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} [t_J - t_{J+k}]^2 = O(l_n) \sum_{a=1}^{q} \sum_{J=1}^{l_n + 2k} O(l_n^{-2}) = O(1).$$

Consequently for all $i \neq j$, $E\left[|B_n(W_i)'| |B_n(W_j)|\right] = O(1)$, where $|B_n(W)|$ be an element wise absolute value.

Also,

$$E\left[b_2^{-1}K_{2ji}^{(1)}(V_d)^2 \mid W_j, S_i\right] = \int b_2^{-1}K_2^{(1)}[b_2^{-1}(V_{dj} - V_{di})]^2 p(V_j; W_j)p(W_j)^{-1}dV_j$$

$$= \int b_2^{-1}K_2^{(1)}(\ )^2 p(V_{di} + b_2\ ; W_j)p(W_j)^{-1}b_2 d$$

$$\sup_{V_d W \in G_{V_d;W}} p(V_d; W) \inf_{W \in G_W} p(W)^{-1} \int K_2^{(1)}(\ )^2 d$$

$$= O(1):$$

Consequently both of the following are true, $E\left[b_2^{-1}K_{2ji}^{(1)}(V_d)^2 \mid W_j, S_i\right] = O(1)$ and $E\left[b_2^{-1}|K_{2ji}^{(1)}(V_d)| \mid W_j, S_i\right] = O(1)$.

Part i.),

$$E\ \|(nh_1^D h_2^{D+1})^{-1}D_d K_{3i}^0 W_d\|$$

Part iii .) Let $\in R^{q(l_n+2k)}$ and consider;

$$\begin{aligned}
&\|n^{-1}B_n^0(M_{dn} - M_{dn}^{l_n})\|_E^2 = E\|n^{-1}B_n^0(M_{dn} - M_{dn}^{l_n})\|_{sp}^2 \\
&= \|n^{-1}\sum_{j=1}^{n} B_n(W_j)[m_d(W_j) - m_d^{l_n}(W_j)]\|_{sp}^2 \\
&= \max n^{-1}\sum_{j=1}^{n}\sum_{g=1}^{n} B_n(W_j)B_n(W_g)^0[m_d(W_j) - m_d^{l_n}(W_j)][m_d(W_g) - m_d^{l_n}(W_g)] \\
&= \max_{\|\|_E=1} n^{-2}\sum_{j=1}^{n}\sum_{g=1}^{n} {}^0B_n(W_j)B_n(W_g)^0\,[m_d(W_j) - m_d^{l_n}(W_j)][m_d(W_g) - m_d^{l_n}(W_g)] \\
&\quad \sup_{W\in G_w}\|m_d(W) - m_d^{l_n}(W)\|^2 \max_{\|\|_E=1} {}^0(Q_{nBB}) \\
&\quad O(l_n^{-2k})\ \max(Q_{nBB})\|\|_E^2 = O(l_n^{-2k})\ \max(Q_{BB}) + o_p(1) = O(l_n^{-2k})
\end{aligned}$$

iii .) follows by taking square roots.

Part iv.):

$$E\|[(n-1)b_2^2]^{-1}K_2^{(1)}(V_d)^0 - {}_{2n}A_n\|I_n(\ i)B_n\|_E^2 = E\|[(n-1)b_2^2]^{-1}\sum_{j\neq i}^{n} B_n(W_j)K_{2ji}^{(1)}(V_d)\ {}_{2j}A_j\|_E^2$$

$$= E\ [(n-1)b_2^2]^{-1}\sum_{j\neq i}^{n}\sum_{g\neq i}^{n} B_n(W_i)^0B_n(W_g)\ {}_{2j}^{d}\ {}_{2g}^{d}K_{2ji}^{(1)}(V_d)K_{2gi}^{(1)}(V_d)A_j\,A_g$$

$$\sup_{X;V\ 2\ G_{X;V}} {}_2^d(X;V)^2\ [(n-1)^2b_2^3]^{-1}\sum_{j\neq i}^{n} E\ B_n(W_j)^0B_n(W_j)E\ b_2^{-1}K_{2ji}^{(1)}(V_d)^2E\ A_j^2\ W_j;V_{dj};S_i\ \|W_j\}$$

$$+[(n-1)^2b_2]^{-2}\sum_{j\neq i}^{n}\sum_{\substack{g\neq i\\ g\neq j}}^{n} E\ \|B_n(W_j)^0\|\|B_n(W_g)\|E\ b_2^{-1}\|K_{2ji}^{(1)}(V_d)\|E\ \|A_j\|\ W_j;V_{dj};S_i;S_g\ \ W_j;S_i;S_g\}^2\}$$

$$O([(n-1)b_2^3]^{-1})E\ B_n(W_j)^0B_n(W_j)E\ b_2^{-1}K_{2ji}^{(1)}(V_d)^2\ W_j\}$$

$$+O([(n-1)^2b_2]^{-2})\sum_{j\neq i}^{n}\sum_{\substack{g\neq i\\ g\neq j}}^{n} E\ \|B_n(W_j)^0\|\|B_n(W_g)\|E\ b_2^{-1}\|K_{2ji}^{(1)}(V_d)\|\ W_j;S_i;S_g\}^2$$

$$= O([(n-1)b_2^3]^{-1})E\ B_n(W_j)^0B_n(W_j)\ + O(b_2^{-2})E\ \|B_n(W_j)^0\|\|B_n(W_g)\|$$

$$= O(l_n[(n-1)b_2^3]^{-1}) + O(b_2^{-2})$$

$$= o(1) + O(b_2^{-2}):$$

$$+ [(n-1)^2 b_2^2]^{-1} \sum_{j \neq i}^{n} \sum_{g \neq i, g \neq j}^{n} E\left[|B_n(W_j)'| |B_n(W_g)| |b_2^{-1} K_{2ji}^{(1)}(V_d)| |b_2^{-1} K_{2gi}^{(1)}(V_d)| |A_j| |A_g| \right]$$

$$= O([(n-1)b_2^3]^{-1}) E\left[B_n(W_j)' B_n(W_j) E\left[b_2^{-1} K_{2ji}^{(1)}(V_d)^2 E\left[A_j^2 \mid W_j, V_{dj}, S_i \right] \mid W_j, S_i \right] \right]$$

$$+ O(b_2^{-2}) E\left[|B_n(W_j)'| |B_n(W_g)| E\left[|b_2^{-1} K_{2ji}^{(1)}(V_d)| E\left[|A_j| \mid W_j, V_{dj}, S_{-j} \right] \mid W_j, S_{-j} \right] \right]^2$$

$$= O([(n-1)b_2^3]^{-1}) E\left[B_n(W_j)' B_n(W_j) E\left[b_2^{-1} K_{2ji}^{(1)}(V_d)^2 \mid W_j, S_i \right] \right]$$

$$+ O(b_2^{-2}) E\left[|B_n(W_j)'| |B_n(W_g)| E\left[|b_2^{-1} K_{2ji}^{(1)}(V_d)| \mid W_j, S_{-j} \right] \right]^2$$

$$= O([(n-1)b_2^3]^{-1}) E\left[B_n(W_j)' B_n(W_j) \right] + O(b_2^{-2}) E\left[|B_n(W_j)'| |B_n(W_g)| \right]$$

$$= O\left(\frac{l_n}{(n-1)b_2^3} \right) + O\left(b_2^{-2} \right) = o(1) + O\left(b_2^{-2} \right)$$

Where the last equality if due to Lemma 3,4.276 O

□

Lemma 6. Let $\{X_i, V_i, A_i\}_{i=1}^n$ be an i.i.d sequence of random variables, where A_i is real valued such that for some $a \geq 2$,

$$\sup_{X_{di} \in G_{X_d}} E[|A_i|^a \mid X_{di}] < C < 1 \quad \text{and} \quad \sup_{V_{di} \in G_{V_d}} E[|A_i|^a \mid V_{di}] < C < 1 :$$

Then under the assumptions of this paper,

$$\sup_{V_{di} \in G_{V_d}} \left[(n-1)b_2^2\right]^{-1} \sum_{j \neq i} \left[K_{2ji}^{(1)}(V_d)^{\frac{d}{2j}} A_j - E\left[K_{2ji}^{(1)}(V_d)^{\frac{d}{2j}} A_j\right]\right] = O_p\left(\left(\frac{\log(n)}{(n-1)b_2^3}\right)^{1/2}\right)$$

$$\sup_{V_{di} \in G_{V_d}} \left[(n-1)b_2\right]^{-1} \sum_{j \neq i} \left[K_{2ji}(V_d)^{\frac{d}{2j}} A_j - E\left[K_{2ji}(V_d)^{\frac{d}{2j}} A_j\right]\right] = O_p\left(\left(\frac{\log(n)}{(n-1)b_2}\right)^{1/2}\right)$$

$$\sup_{X_{di} \in G_{X_d}} \left[(n-1)b_1\right]^{-1} \sum_{j \neq i} \left[K_{1ji}(X_d)^{\frac{d}{1j}} A_j - E\left[K_{1ji}(X_d)^{\frac{d}{1j}} A_j\right]\right] = O_p\left(\left(\frac{\log(n)}{(n-1)b_1}\right)^{1/2}\right) :$$

Proof. Let $m \in \{0, 1\}$ and define $C_s^{(m)}(V_{dj}, V_d) = b_2 K_2^{(m)}[b_2^{-1}(V_{dj} - V_d)]^{\frac{d}{2j}}$ where $V_d \in G_{V_d}$ so that,

$$\begin{aligned} S_n(V_d) &= \left[(n-1)b_2^{m+1}\right]^{-1} \sum_{j \neq i} K_2^{(m)}[b_2^{-1}(V_{dj} - V_d)]^{\frac{d}{2j}} A_j \\ &= \left[(n-1)b_2^{m+2}\right]^{-1} \sum_{j \neq i} b_2 K_2^{(m)}[b_2^{-1}(V_{dj} - V_d)]^{\frac{d}{2j}} A_j \\ &= \left[(n-1)b_2^{m+2}\right]^{-1} \sum_{j \neq i} C_s^{(m)}(V_{dj}, V_d) A_j : \end{aligned}$$

By the Mean Value Theorem, for $V_d^0 \in G_{V_d}$ and some $\in (0, 1)$,

$$\begin{aligned} C_s^{(m)}(V_{dj}, V_d) - C_s^{(m)}(V_{dj}, V_d^0) &= b_2^{\frac{d}{2j}} K_2^{(m)}[b_2^{-1}(V_{dj} - V_d)] - K_2^{(m)}[b_2^{-1}(V_{dj} - V_d^0)] \\ &\quad b_2^{\frac{d}{2j}} K_2^{(m+1)}\left(b_2^{-1}(V_{dj} - V_d) + b_2^{-1}(1-)(V_{dj} - V_d^0)\right) b_2^{-1}(V_d - V_d^0) \\ &\quad \sup_{\in R} K_2^{(m+1)}(\,) \sup_{X,V \in G_{X,V}} {}^{\frac{d}{2}}(X, V) \, V_d - V_d^0 \\ &\quad C \, V_d - V_d^0 : \end{aligned}$$

Let $\{B_n\}_{n=1}^1$ be a nondecreasing sequence of real positive numbers s.t. $\sum_{n=1}^1 B_n^{-a} < 1$ and define,

$$S_n(V_d) = \left[(n-1)b_2^{m+1}\right]^{-1} \sum_{j \neq i} C_s^{(m)}(V_{dj}, V_d) A_j \, 1\{|A_j| \leq B_n\}:$$

where $1\{\,\}$ is an indicator function. Consider,

$$\sup_{V_d \in G_{V_d}} S_n(V_d) - E(S_n(V_d)) \qquad \sup_{V_d \in G_{V_d}} S_n(V_d) - \|$$

T_{32}:

$$\sup_{V_d \in G_{V_d}} |E\,S(V_d^l) - S(V_d)| \le \sup_{V_d \in G_{V_d}} [(n-1)b_2^{m+2}]^{-1} \sum_{j \neq i} E\,|C_s^{(m)}(V_{d_j}; V_d^l) - C_s^{(m)}(V_{d_j}; V_d)|\,A_j\,1\{|A_j| > B_n\}$$

$$\le \sup_{V_d \in G_{V_d}} [(n-1)b_2^{m+2}]^{-1} \sum_{j \neq i} E\,|C(V_d^l, V_d)|\,A_j\,1\{|A_j| > B_n\}$$

$$= O\left(\frac{n}{b_2^{m+2}}\right)(n-1)^{-1} \sum_{j \neq i} E(|A_j|) = O\left(\frac{n}{b_2^{m+2}}\right):$$

T_{33}: Let $\{\varepsilon_n\}_{n=1}^{\infty}$ be a sequence of positive real numbers s.t. $\lim_{n\to\infty} \varepsilon_n = 0$, and note that,

$$P\left(\max_{1 \le l \le n} |S(V_d^l) - E\,S(V_d^l)| > \varepsilon_n\right) \le P\left(\bigcup_{l=1}^{n} |S(V_d^l) - E\,S(V_d^l)| > \varepsilon_n\right)$$

$$\le \sum_{l=1}^{n} P(|S(V_d^l) - E\,S(V_d^l)| > \varepsilon_n) \le n\varepsilon_n$$

E S

Consequently,

$$P \; jS \; (V_d) \quad E \; S \; (V_d^{|}) \; j \quad "_n$$

Consequently, by Markov's inequality,

$$T_{33} = \max_{i\ l_n} jS\ (V_d^l) \quad E$$

$$E\ b_1^{-1}K_{1ji}(X_d)\ \tfrac{d}{1j}H(Z_{ci};X_{di})jS_i$$

T_1 T_2 T_3:

$$T_1 = \underset{Z}{E}\ b_1^{-1}K_{1ji}(X_d)\ \tfrac{d}{1j}E[Z_{cj}jX_j;V_j]jS_i]$$

$$= \int b_1^{-1}K_{1ji}(X_d)\frac{g(X_{dj};V_j)}{p(X_j;V_j)}\int Z_{cj}\frac{p(Z_{cj};X_j;V_j)}{p(X_j;V_j)}dZ_{cj}\,p($$

$$= \int b_1^{-1}K_{1ji}(X_d)\int\frac{g(X_j;V_j)}{p(X_j;V_j)}Z_{cj}\frac{p(Z_{cj};X_j;V_j)}{p(X_{dj})}dZ_{cj}\,($$

$$= \int b_1^{-1}K_{1ji}(X_d)\ H(Z_{cj};X_{dj})\quad H(Z_{ci};X_{di}) + H(\quad$$

$$= \int b_1^{-1}K_{1ji}(X_d)\ H(Z_{cj};X_{dj})\quad H(Z_{ci};X_{di})\ dX_{dj} + H(Z_{ci};X_{di})\quad b_1^{-1}K_{1ji}(X_d)dX_{dj}$$

$$= O(b_1^{-1}) + H(Z_{ci};X_{di}):$$

The last equality is due to Assumption A4 and i.), T_2: By Assumption A4 and i.),

$$T_2 = E\ b_1^{-1}K_{1ji}(X_d)\ \tfrac{d}{1j}\ H(Z_{cj};X_{dj})\quad H(Z_{ci};X_{di})\ jS_i\ = O(b_1^{-1})$$

T_3:

$$E\ b_1^{-1}K_{1ji}(X_d)\ \tfrac{d}{1j}H(Z_{ci};X_{di})jS_i\ = H(Z_{ci};X_{di})E\ b_1^{-1}K_{1ji}($$

$$= H(Z_{ci};X_{di})b_1^{-1}\quad K\quad ;V_j)dX_j\,dV_j$$

$$= H(Z_{ci};X_{di})b_1^{-1}\quad H(Z_{ci};X_{di}):$$

In summary,

$$E\ b_1^{-1}K_{1ji}(X_d)\ \tfrac{d}{1j}\ \tfrac{d}{1ji}\ jS_i\ = T_1\quad T_2\quad T_3 = O(b_1^{-1}) +\quad)\quad H(Z_{ci};X_{di}) = O(b_1^{-1}):$$

Part iv.): The proof of iv:) follows, mutatis mutandis, iii).

□

Lemma 8. Under the assumptions A1 - A6,

i) $E\ \ _i\ _{ci}H_2^{(1)\,d}(Z_{mi})B_L(W_{ai})\ =$

ii) $E\ \ _i\ _{ci}H_2^{(1)\,d}(Z_{mi})V_{di}B_L(W_{ai})B_J($

iii) $E\ \ \tfrac{2}{1}\ \ z_{ci};X_{di})\quad B_L\quad W\quad O(ll_3$)

ii))

$$B_a(L;I_n) = W_a(A_a(L;I_n)) \quad R$$
$$C_a(L;I_n) = (A_a(L;I_n);X;V;W_a) \quad R^{2D+1}$$
$$D_{ac}(L;I_n) = (A_a(L;I_n);Z_c;X;V;W_a) \quad R^{2D+2}$$
$$E(L;I_n) = (A_a(L;I_n);X;V) \quad R^{2D}$$
$$F_c(L;I_n) = (A_a(L;I_n);Z_c;X;V) \quad R^{2D+1}:$$

Note that $C_a(L;I_n) \quad E(L;I_n) \quad B_a(L;I_n)$, and $D_{ac}(L;I_n) \quad F_c(L;I_n) \quad B_a(L;I_n)$. As a result for any functions $f(Z_c;X;V;W_a) : R^{2D+2} \; ! \; R^+$, and $f(X;V;W_a) : R^{2D+1} \; ! \; R^+$,

$$\int_{D_{ac}(L;I_n)} f(Z_c;X;V;W_a)p(Z_c;X;V;W_a)d(Z_c;X;V;W_a)$$
$$\int_{F_c(L;I_n) \quad B_a(L;I_n)} f(Z_c;X;V;W_a)p(Z_c;X;V;W_a)d(Z_c;X;V;W_a):$$

and

$$\int_{C_a(L;I_n)} f(X;V;W_a)p(X;V;W_a)d(X;V;W_a)$$
$$\int_{E(L;I_n) \quad B_a(L;I_n)} f(X;V;W_a)p(X;V;W_a)d(X;V;W_a):$$

Furthermore by the remarks to assumption A1,

$$\int_{B_a(L;I_n)} dW_a = W_a \Big|_{t_L}^{t_{L+k}} = (t_{L+k} \quad t_L) \quad CkI_n^{\,1}:$$

Consider,

$$P\{B_L(W_{ai}) > 0\} = P\{W_a \, 2 \, B_a(L;I_n)\} = \int_{B_a(L;I_n)} p(W_a)dW_a = P\{f! \; 2 \; : \; W_a(!) \, 2 \, B_a(L;I_n)g$$
$$= P(A_a(L;I_n)) = P\{f! \; 2 \; : [Z_c(!) \; X(!)^0 V(!)^0]^0 2 \, F_c(L;I_n)g$$
$$= P\{F_c(L;I_n)\} = \int_{F_c(L;I_n)} p(Z_c;X;V)d(Z_c;X;V):$$

Similarly,

$$P\{B_L(W_{ai}) > 0\} = P\{A_a(L;I_n)\} = P\{f! \; 2 \; : [X(!)^0 V(!)^0]^0 2 \, E(L;I_n)g$$
$$= P\{E(L;I_n))\} = \int_{E(L;I_n)} p(X;V)dXdV:$$

Consequently,

$$P\{B_L(W_a) > 0) = \int_{B_a(L;I_n)} p(W_a)dW_a = \int_{F_c(L;I_n)} p(Z_c;X;V)dZ_cdXdV = \int_{E(L;I_n)} p(X;V)dX;dV:$$

Now consider,

$$\int_{C_a(L;I_n)} p(W_a)dW_a \quad \sup_{W_a 2 G_{W_a}} p(W_a) \int_{C_a(L;I_n)} dW_a = CkI_n^{\,1} = O(I_n^{\,1}):$$

Consequently,

$$0 \quad \int_{F_c(L;I_n)} p(Z_c;X;V)dZ_cdXdV = \int_{E(L;I_n)} p(X;V)dX;dV = \int_{B_a(L;I_n)} p(W_a)dW_a = O(I_n^{\,1}):$$

Hereafter notation is simpli ed in the following way, $B_a \quad B_a(L; I_n)$, $C_a \quad C_a(L; I_n)$, $D_{ac} \quad D_{ac}(L; I_n)$, $E \quad E(L; I_n)$, and $F_c \quad F_c(L; I_n)$.

Part i.):
Case 1: $Z_c \notin W_a$,

$$E \; {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi}) B_L(W_{ai}) \qquad E \; {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi})\; B_L(W_{ai}) 1\; B_L(W_{ai}) > 0$$

$$= \int_{D_{ac}} {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi})\; B_L(W_{ai}) p(Z_{ci}; X_i; V_i; W_{ai}) dZ_{ci}\, dX_i\, dV_i\, dW_{ai}$$

$$\int_{F_c \quad B_a} {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi})\; B_L(W_{ai}) p(Z_{ci}; X_i; V_i; W_{ai}) dZ_{ci}\, dX_i\, dV_i\, dW_{ai}$$

$$= \int_{F_c} {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi}) \int_{B_a} B_L(W_{ai}) p(Z_{ci} X_i; V_i; W_{ai}) p(Z_{ci}; X_i; V_i)\; {}^{1} dW_{ai}\, p(Z_{ci}; X_i; V_i) dZ_{ci}\, dX_i\, dV_i$$

$$jj b_L(W_{ai}) jj_2 \; {}^{1} \sup_{W_a 2 G_{W_a}} b_L(W_a) \int_{Z_c;X;V \; 2 G_{Z_c;X;V;W_a}} \sup p(Z_c; X; V; W_a) \int_{Z_c;X;V \; 2 G_{Z_c;X;V}} \inf p(Z_c; X; V)\; {}^{1}$$

$$\int_{F_c} {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi}) \int_{B_a} dW_{ai}\; p(Z_{ci}; X_i; V_i) dZ_{ci}\, dX_i\, dV_i$$

$$O\; I_n {}^{1=2} \sup_{X;V \; 2 G_{XV}} (X; V)\; {}_c H_2^{(1)\,d}(Z_m) \int_{F_c} p(Z_{ci}; X_i; V_i) dZ_{ci}\, dX_i\, dV_i$$

$$= O(I_n {}^{3=2}):$$

Case 2: $Z_c = W_a$,

$$E \; {}_i\, {}_{ci} H_2^{(1)\,d}(Z_{mi}) B_L(W_{ai}) \qquad E \; {}_i\, W_{ai} \quad H\ (W_{ai})\; H_2^{(1)\,d}(Z_{mi})\; B_L(W_{ai})$$

$$\int {}_i\, H_2^{(1)\,d}(Z_{mi})\; W_{ai}\; B_L(W_{ai}) p(X_i; V_i; W_{ai}) dX_i\, dV_i\, dW_{ai}$$

$$C_a$$

$$+ \int_{C_a} {}_i\, H_2^{(1)\,d}(Z_{mi})\; H\ (W_{ai})\; B_L(W_{ai}) p(X_i; V_i; W_{ai}) dX_i\, dV_i\, dW_{ai}$$

$$\int {}_i\, H_2^{(1)\,d}(Z_{mi}) \int W_{ai}\; B_L(W_{ai}) p(X_i; V_i; W_{ai}) p(X_i; V_i)\; {}^{1} dW_{ai}\; p(X_i; V_i) dX_i\, dV_i$$

$$E \qquad B_a$$

$$+ \int_{E} {}_i\, H_2^{(1)\,d}(Z_{mi})\; H\ (W_{ai}) \int_{B_a} B_L(W_{ai}) p(X_i; V_i; W_{ai}) p(X_i; V_i)\; {}^{1} dW_{ai}\; p(X_i; V_i) dX_i\, dV_i$$

$$jj b_L(W_{ai}) jj_2 \; {}^{1} \sup_{W_a 2 G_{W_a}} b_L(W_a) \sup_{X;V \; 2 G_{X;V;W_a}} p(X; V; W_a) jW_a j \inf_{X;V \; 2 G_{X;V}} p(X; V)\; {}^{1}$$

$$i \; \Big(\int \sup_{X;V \; 2 G_i} {}_i\, H_2^{(1)\,d}(Z_{mi})\; 1 + 1\; H\ (W_{ai}) B_L(W_{ai}\; 1 dW_{ai}\; p(X_i; V_i) dX_i\, dV_i \Big)$$

$$E \qquad E \; (X_{ai} H_2^{(1)\,d}(Z_m) \qquad B_a$$

1a9/F11 9.9626 Tf 3.875 0 Td [(W)]TJ/F10 6.9738 Tf 9.40-1.494 Td [(mi)]TJ

$$O(l_n^{1=4})E \quad {}^{2}_{1}\,{}^{2}_{ci}H_2^{(1)\,d}(Z_{mi})^2 B_L(W_{ai}) \quad {}^{1=2}:$$

Note that a careful inspection of the proof of Part (i), yields the following conclusion,

$$E \quad {}^{2}_{1}\,{}^{2}_{ci}H_2^{(1)\,d}(Z_{mi})^2 B_L(W_{ai}) \quad = O(l_n{}^{3=2}):$$

Consequently, $E \quad {}^{2}_{1}\,{}^{2}_{ci}H_2^{(1)\,d}(Z_{mi})^2 B_L(W_{ai})^2 \quad {}^{1=2} = O(l_n^{1=4})O(l_n{}^{3=4}) = o(l_n{}^{1=2})$.

Part iv.):

$$E \quad {}^{2}_{i}\,{}^{2}_{ci}H_2^{(1)\,d}(Z_{mi})^2 V_{di}^2 B_L(W_{ai})^2 B_J(W_{li})^2 \quad {}^{1=2}$$

$$jjb_L(W_{li})jj_2 \quad {}^{1}\; jjb_L(W_{ai})jj_2 \quad {}^{1=2} \sup_{W_a;W_l 2 G_{W_a;W_l}} b_L(W_l)b_L(W_a)^{1=2}E \quad {}^{2}_{1}\,{}^{2}_{ci}H_2^{(1)\,d}(Z_{mi})^2 V_{di}^2 B_L(W_{ai}) \quad {}^{1=2}$$

$$= O(l_n^{1=2})O(l_n^{1=4}) \sup_{V_d 2 G_{V_d}} jV_d jE \quad {}^{2}_{1}\,{}^{2}_{ci}H_2^{(1)\,d}(Z_{mi})^2 B_L(W_{ai}) \quad {}^{1=2} = O(l_n^{3=4})O(l_n{}^{3=4}):$$

□

Lemma 9. Suppose that for some sequence of positive real numbers $fT_n g_{n=1}^{1}$,

$$\sup_{X_{di} 2 G_X} j\hat{p}(X_{di}) \quad p(X_{di})j = O_p(T_n); \quad \text{and} \quad \sup_{V_{di} 2 G_V} j\hat{p}(V_{di}) \quad p(V_{di})j = O_p(T_n):$$

Then, under the assumptions A1 - A6,

$$\hat{g}(X_i;\hat{V}_i) \quad g(X_i;V_i) = 2 \sum_{d=1}^{p} g(X_{\quad di};V_i)\,\hat{p}(X_{di}) \quad p(X_{di})$$

$$= \hat{p}(T_j)\left[\prod_{d=j+1}^{D} p(T_d)\right] + \left[\hat{p}(T_j) - p(T_j)\right]\prod_{d=j+1}^{D} p(T_d)$$
$$= \hat{p}(T_j) - p(T_j) + p(T_j)\, A_{j+1} + C(j; f_j g)$$
$$= B_j A_{j+1} + C(j; f_j g).$$

Consequently,

$$\begin{aligned} A_1 &= B_1 A_2 + C(1; f_1 g) \\ &= B_1 B_2 A_3 + C(2; f_2 g) + C(1; f_1 g) \\ &= B_1 B_2 A_3 + B_1 C(2; f_2 g) + C(1; f_1 g) \\ &= B_1 B_2 B_3 A_4 + C(3; f_3 g) + B_1 C(2; f_2 g) + C(1; f_1 g) \\ &= B_1 B_2 B_3 A_4 + B_1 B_2 C(3; f_3 g) + B_1 C(2; f_2 g) + C(1 f_1 g) \\ &\vdots \qquad\qquad \vdots \qquad\qquad \vdots \\ &= A_D \prod_{i=1}^{D-1} B_i + \sum_{j=1}^{D-1} C(j+1; f_{j+1} g) \prod_{1 \le i \le j} B_i + C(1; f_1 g). \end{aligned}$$

Now note that,

$$A_D \prod_{i=1}^{D-1} B_i = [\hat{p}(T_1) - p(T_1)] + p(T_1)\,[\hat{p}(T_2) - p(T_2)] + p(T_2)$$
$$\hat{p}(T_{D-1}) - p(T_{D-1}) + p(T_{D-1})\,\hat{p}(T_D) - p(T_D)$$

If one expands this product into a sum consisting solely of terms of the form $D(\ _{j+1})$ then, for each $j \in \{1, 2, \ldots, D-1\}$, there are exactly $\binom{D-1}{j}$ of these terms within this sum. Furthermore, for all $_i$ one has

$$D(\ _i) = \left[\prod_{d=i}^{D} p(T_d)\right]\left[\prod_{d=i}^{D} \hat{p}(T_d) - p(T_d)\right]$$
$$\left[\prod_{d=i}^{d2_i} \sup_{T_d \in G_{T_d}} p(T_d)\right]\left[\prod_{d=i}^{d2_i} \sup_{T_d \in G_{T_d}} \hat{p}(T_d) - p(T_d)\right]$$
$$= O(1) O_p\left(L_n^{\#(\ _i)}\right) = O_P\left(L_n^i\right).$$

Consequently one can write

$$A_D \prod_{i=1}^{D-1} B_i = \sum_{j=1}^{D} D(f_j g) + \sum_{j=1}^{D-1} \binom{D-1}{j} O_p(L_n^{j+1}) = \sum_{j=1}^{D} D(f_j g) + O_p(L_n^2).$$

Similarly,

$$C(j+1; f_{j+1} g) \prod_{1 \le i \le j} B_i = [\hat{p}(T_1) - p(T_1)] + p(T_1)\,[\hat{p}(T_2) - p(T_2)] + p(T_2)$$
$$[\hat{p}(T_j) - p(T_j)] + p(T_j)\left[\hat{p}(T_{j+1}) - p(T_{j+1})\right]\prod_{j+2 \le d} p(T_d).$$

Furthermore, if one where to expand this product into a sum consisting soley of terms of the form $D(\ _{m+1})$ then for each $m \in \{1, 2, \ldots, j\}$, there are exactly, $\binom{j}{m}$ of these in the sum. Consequently, one can write,

$$C(j+1; f_{j+1} g) \prod_{1 \le i \le j} B_i = D(f_{j+1} g) + \sum_{m=1}^{j} \binom{j}{m} O_p(L_n^{m+1}) = D(f_{j+1} g) + O_p(L_n^2).$$

As a result, one has,

$$\sum_{j=1}^{\infty} C(j+1;\{j+1\}) \prod_{1 \le i \le j} B_i = \sum_{j=1}^{\infty} D(\{j+1\}) + O_p(L_n^2):$$

Lastly note that $C(1;\{1\}) = D(\{1\})$ so that,

$$\begin{aligned} A_1 &= A_D \prod_{i=1}^{\infty} B_i + \sum_{j=1}^{\infty} C(j+1;\{j+1\}) \prod_{1 \le i \le j} B_i + C(1;\{1\}) \\ &= \sum_{j=1}^{D} D(\{j\}) + O_p(L_n^2) + \sum_{j=1}^{\infty} D(\{j+1\}) + O_p(L_n^2) + D(\{1\}) \\ &= 2 \sum_{j=1}^{D} D(\{j\}) + O_p(L_n^2): \end{aligned}$$

□

Lemma 10. Under the assumptions A1 - A6, $n^{-1} \sum_{i=1}^{n} \psi_i u_i(\hat{\theta}_i - \theta_i) = o_p(n^{-1/2})$.

Proof.

$\hat{p}(X_i;\hat{V}_i) - p(X_i;V_i) = [\hat{p}(X_i;\hat{V}_i)p(X_i;V_i)]^{-1} p(X_i;V_i)$.8593 6.9738 Tf 5.98 3.615 Td [( )]TJ/F7 6.9738 Tf 6.227 0 Td [(1

$$= 2\sum_{d=1}^{p} p(X_i;V_i)\ {}^{1}g(X_{di};V_i)\ \hat{p}(X_{di}) \quad p(X_{di})$$

$$+ 2\sum_{d=1}^{p} p(X_i;V_i)\ {}^{1}g(X_i;V_{di})\ \hat{p}(\hat{V}_{di}) \quad p(V_{di}) \ + O_p(L_{0n}^2)$$

$$= 2\sum_{d=1}^{p} p(X_{di})\ {}^{1}{}_{i}\ \hat{p}(X_{di}) \quad E(\hat{p}(X_{di})\ X_{di}) \ + 2\sum_{d=1}^{p} p(X_{di})\ {}^{1}{}_{i}[E(\hat{p}(X_{di})jX_{di}) \quad p(X_{di})]$$

$$+ 2\sum_{d=1}^{p} p(V_{di})\ {}^{1}{}_{i}\ \hat{p}(\hat{V}_{di}) \quad \hat{p}(V_{di}) \ + 2\sum_{d=1}^{p} p(V_{di})\ {}^{1}{}_{i}\ \hat{p}(V_{di}) \quad E(\hat{p}(V_{di})jV_{di})$$

$$+ 2\sum_{d=1}^{p} p(V_{di})\ {}^{1}{}_{i}\ E(\hat{p}(V_{di})jV_{di}) \quad p(V_{di}) \ + o_p(n^{\ 1=2})$$

$$2\sum_{d=1}^{p} T_{1d} + 2\sum_{d=1}^{p} T_{2d} + 2\sum_{d=1}^{p} T_{3d} + 2\sum_{d=1}^{p} T_{4d} + 2\sum_{d=1}^{p} T_{5d} + o_p(n^{\ 1=2}):$$

Consider also,

$$p(X_i;V_i)\ {}^{1}{}_{i}\ p(X_i;V_i) \quad \hat{p}(X_i;\hat{V}_i) = (\quad 1)p(X_i;V_i)\ {}^{1}{}_{i}\ \hat{p}(X_i;\hat{V}_i) \quad \hat{p}(X_i;V_i)$$

$$p(X_i;V_i)\ {}^{1}{}_{i}\ \hat{p}(X_i;V_i) \quad E(\hat{p}(X_i;V_i)jX_i;V_i)$$

$$p(X_i;V_i)\ {}^{1}{}_{i}\ E(\hat{p}(X_i;V_i)jX_i;V_i \quad p(X_i;V_i)$$

$$T_6 \quad T_7 \quad T_8:$$

where, $\overset{(2)}{1}{}^{d}(i;j\ ;c) = \overset{d}{1}(i;j\ ;c) + \overset{d}{1}(j;i\ ;c)$

$E\ \overset{d}{1}(i;j\ ;c) = E_{ci}\, p(X_{di})$